

QM016/1
Mathematics
Paper 1
Semester I
2006/2007
2 hours

QM016/1
Matematik
Kertas 1
Semester I
2006/2007
2 jam



BAHAGIAN MATRIKULASI
KEMENTERIAN PELAJARAN MALAYSIA
MATRICULATION DIVISION
MINISTRY OF EDUCATION MALAYSIA

PEPERIKSAAN SEMESTER PROGRAM MATRIKULASI
MATRICULATION PROGRAMME EXAMINATION

MATEMATIK
Kertas 1
2 jam

JANGAN BUKA KERTAS SOALAN INI SEHINGGA DIBERITAHU.
DO NOT OPEN THIS BOOKLET UNTIL YOU ARE TOLD TO DO SO.

CHOW CHOON WOOI

Kertas soalan ini mengandungi **13** halaman bercetak.
This booklet consists of 13 printed pages.

© Bahagian Matrikulasi

INSTRUCTIONS TO CANDIDATE:

This question booklet consists of **10** questions.

Answer **all** questions.

The full marks allocated for each question or section is shown in the bracket at the end of each question or section.

All steps must be shown clearly.

Only non-programmable scientific calculator can be used.

Numerical answers can be given in the form of π , e , surd, fractions or up to three significant figures, where appropriate, unless stated otherwise in the question.

LIST OF MATHEMATICAL FORMULAE

Arithmetic Series:

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d]$$

Geometric Series:

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} \quad \text{for } r < 1$$

Binomial Expansion:

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n, \text{ where } n \in \mathbb{N} \text{ and}$$

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \text{ for } |x| < 1$$

1. If $P = \begin{bmatrix} 1 & -2 \\ 1 & 1 \\ 0 & -1 \end{bmatrix}$ and $Q = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix}$, find matrix R such that

$$R + 2(PQ) = \begin{bmatrix} 0 & 2 & 2 \\ -2 & 4 & 3 \\ -4 & 5 & 3 \end{bmatrix}.$$

[5 marks]

2. By substituting $a = 2^x$, solve the equation

$$4^x + 3 = 2^{x+2}.$$

[6 marks]

3. Obtain the solution set for

$$|2x+1| > -x^2 + 4.$$

[7 marks]

4. The sum of the first k terms of an arithmetic series is 777. The first term is -3 and the k -th term is 77. Obtain the value of k and the eleventh term of the series.

[7 marks]

5. (a) Find the values of A , B , C and D for the expression $\frac{4x^3 - 3x^2 + 6x - 27}{x^4 + 9x^2}$

in the form of partial fractions $\frac{A}{x} + \frac{B}{x^2} + \frac{Cx+D}{x^2+9}$ where A , B , C and D

are constants.

[5 marks]

- (b) Given $A = \begin{bmatrix} -2 & 0 & 0 \\ -4 & 6 & -2 \\ 6 & -4 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 & 0 \\ -1 & \frac{1}{5} & -\frac{1}{5} \\ -1 & -\frac{2}{5} & -\frac{3}{5} \end{bmatrix}$. Show that $AB = kI$

where k is a constant and I is an identity matrix. Find the value of k and hence obtain A^{-1} .

[5 marks]

6. (a) Given the complex number z and its conjugate \bar{z} satisfy the equation $z\bar{z} + 2z = 12 + 6i$. Find the possible values of z .

[6 marks]

- (b) An equation in a complex number system is given by

$$z = \frac{1}{(z_1 - z_2)} + \frac{1}{\bar{z}_1}$$

where $z_1 = 1 + 2i$ and $z_2 = 2 - i$. Find

- (i) the value of z in the Cartesian form $a + ib$

[3 marks]

- (ii) the modulus and argument of z .

[3 marks]

7. (a) Find the solution set of the inequality

$$\frac{1}{3 - 2x} < \frac{1}{x + 4}$$

[5 marks]

- (b) Solve the following inequality equation for all x is real numbers. Write your answer in set form.

$$4 - \left| \frac{3 - 2x}{1 + x} \right| \geq 1$$

[7 marks]

8. (a) Show that $(x - 3)$ is a factor of the polynomial

$$P(x) = x^3 - 2x^2 - 5x + 6.$$

Hence, factorize $P(x)$ completely.

[4 marks]

- (b) If $f(x) = ax^2 + bx + c$ leaves remainder 1, 25 and 1 on division by $(x - 1)$, $(x + 1)$ and $(x - 2)$ respectively, find the values of a , b and c . Hence, show that $f(x)$ has two equal real roots.

[9 marks]

9. (a) Find the first four terms in the binomial expansion of the following functions:

(i) $\sqrt{1+2x}$ [2 marks]

(ii) $\frac{1}{(1-x)^2}$. [2 marks]

(b) Hence, expand $\sqrt{\frac{1+2x}{(1-x)^4}}$ in ascending power of x up to the term containing x^3 . By putting $x = \frac{1}{10}$, show that $\sqrt{12000}$ is approximately $\frac{10935}{100}$.

[9 marks]

10. A doctor prescribed to a patient 13 units of vitamin A, 22 units of vitamin D and 31 units of vitamin E each day. The patient can choose from the combination of three brands of capsules; L, M and N. Each capsule of brand L contains 1 unit each of vitamins A, D and E. Each capsule of brand M contains 1 unit of vitamins A, 2 units of vitamin D, and 3 units of vitamin E. Each capsule of brand N contains 4 units of vitamins A, 7 units of vitamin D and 10 units of vitamin E. The above information is summarized in the following table:

Type of Vitamins	Brand of Capsules			Total Unit of Vitamins
	L	M	N	
A	1	1	4	13
D	1	2	7	22
E	1	3	10	31

By using x as the number of capsules of brand L, y the number of capsules of brand M and z the number of capsules of brand N,

(a) form a system of linear equations from the above information. [2 marks]

(b) write the above system of linear equations in the form of matrix equation: $AX = B$, where A is the coefficient matrix, X is the variable matrix, and B is the constant matrix. Solve the system of equations by using the Gauss-Jordan elimination method. [8 marks]

(c) determine the possible combinations of the number of capsules of brand L, M and N to be taken each day. [3 marks]

- (d) If brand L costs 10 cents per capsule, brand M costs 30 cents per capsule and brand N costs 60 cents per capsule. Determine the combination that will minimize the patient's daily cost.

[2 marks]

END OF QUESTION BOOKLET